

Apr. 1994

A trial to find an elliptic quantum algebra for sl_2 using the Heisenberg and Clifford algebra

Jun'ichi Shiraishi

*Department of physics, University of Tokyo
Bunkyo-ku, Tokyo 113, Japan*

ABSTRACT

A Heisenberg-Clifford realization of a deformed $U(sl_2)$ by two parameters p and q is discussed. The commutation relations for this deformed algebra have interesting connection with the theta functions.

Introduction Recently it is found in the paper [1] that the so called $RLL = LLR$ formalism successfully leads an elliptic quantum algebra for $\widehat{sl}_2$ which enables us to treat the eight-vertex model by using the infinit dimensional algebra technique as we could do for the six-vertex model (see references in [1]).

At this stage it seems very hard to work directly with the elliptic algebra mainly because we do not have its free field realization. The commutation relations among the L operators for the elliptic algebra are so complicated that it's not an easy task to find the right way to bosonize the algebra. So, we need to start from a much simple toy model and study its bosonization.

We will discuss in the following about an two parameter deformation of the algebra $U(sl_2)$ using the Heisenberg algebra $[\partial_z, z] = 1$ and the Clifford algebra $\mathcal{O}_{2n+1}\mathcal{O}_{2n+1} = 1$, $\mathcal{O}_{2m+1}\mathcal{O}_{2n+1} = -\mathcal{O}_{2n+1}\mathcal{O}_{2m+1}$ for $m, n \in \mathbf{Z}_{\geq 0}$ and $m \neq n$.

an attempt at obtaining an elliptic sl_2 algebra using the Heisenberg algebra

As a toy model for the elliptic algebra introduced in [1], we want to discuss the following rather simple commutation relation. Let us introduce two parameters p, q and define the p, q deformation of the universal enveloping algebra for sl_2 as an associative algebra generated by e, f , and $t^{\pm 1}$ having the following commutation relations

relations:

$$\begin{aligned} tet^{-1} &= q^2 e, & tft^{-1} &= q^{-2} f, \\ [e, f] &= \sum_{n \in \mathbf{Z}} (-1)^n t^{2n+1} p^{(n+1/2)^2}. \quad \dots (*) \end{aligned}$$

We note that the form of the right hand side of the equation (*) is nothing but the theta function.

It is well known that the Lie algebra sl_2 and the quantum algebra $U_q(sl_2)$ have the following Heisenberg realization [2]

sl_2 case:

$$e = -\partial_z, \quad h = -2z\partial_z + \lambda, \quad f = z^2\partial_z - \lambda z,$$

$U_q(sl_2)$ case:

$$e = -\frac{1}{z} [\vartheta], \quad t = q^{-2\vartheta+\lambda}, \quad f = z [\vartheta - \lambda],$$

here, we set $\vartheta = z\partial_z$ and $[A] = (q^A - q^{-A})/(q - q^{-1})$.

Now we state our main result.

Proposition: *We have the following Heisenberg-Clifford realization for our elliptic sl_2 algebra,*

$$\begin{aligned}
e &= -\frac{1}{z} \sum_{n \in \mathbf{Z}} a_{2n+1} q^{(2n+1)\vartheta} p^{(n+1/2)^2/2} \mathcal{O}_{|2n+1|}, \\
t &= q^{-2\vartheta+\lambda}, \\
f &= z \sum_{n \in \mathbf{Z}} b_{2n+1} q^{(2n+1)(\vartheta-\lambda)} p^{(n+1/2)^2/2} \mathcal{O}_{|2n+1|},
\end{aligned}$$

where

$$\begin{aligned}
a_{2n+1} &= \frac{q^{-(2n+1)(\lambda+1)/2} (-1)^{-n/2}}{\sqrt{q^{2n+1} - q^{-2n-1}}}, \\
b_{2n+1} &= \frac{q^{(2n+1)(\lambda+1)/2} (-1)^{-n/2}}{\sqrt{q^{2n+1} - q^{-2n-1}}}
\end{aligned}$$

for $n \in \mathbf{Z}$.

Proof: Let $n, m \in \mathbf{Z}$ and $n \neq m$. To satisfy the commutation relation $(*)$ we must have

$$a_{2n+1} b_{2n+1} = \frac{(-1)^{-n}}{q^{2n+1} - q^{-2n-1}},$$

$$a_{2n+1} b_{2m+1} q^{(n-m)(\lambda+1)} - a_{2m+1} b_{2n+1} q^{-(n-m)(\lambda+1)} = 0.$$

These equations can be solved and have the solution stated as above. *QED*

We discussed an Heisenberg realization of a two parameter deformation of the universal enveloping algebra of sl_2 . So far, we have not found a Hopf algebra structure which is necessary to construct tensor product representations of our algebra. The connection between our algebra and the algebra defined in [1] is not clear. We hope that to do a loop-algebraic extension of our algebra is also possible in the similar manner as discussed in [2]. It is an interesting problem to think why the operators e and f resemble theta functions except for the q -dependent factor when we bosonize them. How can we interpret the infinite dimensional Clifford algebra that is so trivial that we did not need when $p = 0$?

Acknowledgements The author would like to thank H. Awata, T. Eguchi, O. Foda, M. Jimbo, T. Miwa and K. Sugiyama for helpful discussions. He is grateful to kind hospitality at YITP and RIMS.

References

- [1] Foda O, Iohara K, Jimbo M, Kedem R, Miwa T and Yan H, An elliptic quantum algebra for $\widehat{sl}_2$, *RIMS preprint* (1994); Notes on highest weight modules of the elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}_2)$, *RIMS preprint* (1994).
- [2] Awata H, Odake S, Shiraishi J, Free boson realization of $U_q(\widehat{sl}_N)$, *to appear in Comm. Math. Phys.*, (1993).